

Free Questions for FPGEE by vceexamstest

Shared by Morin on 29-01-2024

For More Free Questions and Preparation Resources

Check the Links on Last Page

Question Type: MultipleChoice

Drugs shown to be effective in the prevention of Travellers'

Diarrhea, which might have been prescribed for SM include:

I doxycycline

II cotrimoxazole

III amoxicillin

Options:

- A- I only
- B- III only
- C- I and II only
- D- II and III only
- E- I, II and III

Answer:

Question Type: MultipleChoice

Organisms commonly implicated in the cause of Traveller's Diarrhea include:

I Bacteroides fragilis

II Escherichia coli

III Shigella sp

Options:

- A- I only
- B- III only
- C- I and II only
- D- II and III only
- E- I, II and III

Answer:	
Question 3	
Question Type: MultipleChoice	
Which of the following drugs could be the cause of SM's complaints?	
Options:	
A- Bismuth subsalicylate	
B- Cotrimoxazole	
C- Doxycycline	
D- Amoxicillin	
E- Loperamide	
Answer:	
A	

Question Type: MultipleChoice

The following prescription is received in a community pharmacy:

Losec® 20 mg

S: Take 1 tab BID

M: 2 weeks

Which of the following is correct labelling for this prescription?

Options:

- A- Take one tablet twice weekly (28 tablets).
- B- Take one tablet twice weekly (4 tablets).
- C- Take two tablets twice daily (56 tablets).
- D- Take one tablet twice daily (28 tablets).
- E- Take one tablet twice daily (14 tablets)

Answer:

Question Type: MultipleChoice

Dopamine is useful in the treatment of cardiogenic shock because it:

I selectively dilates renal and mesenteric vascular beds.

Il does not induce peripheral vasoconstriction.

III decreases the force of myocardial contraction.

Options:

- A- I only
- B- III only
- C- I and II only
- D- II and III only
- E- I, II and III

Answer:

Α

Question 6

Question Type: MultipleChoice

Following the ethical principle of nonmaleficence requires that pharmacists:

Options:

- A- respect the rights of patients to make choices
- B- do good to patients, placing the benefit of the patient above all else
- C- avoid, remove or prevent harm from people
- D- act with fairness, allowing people access to pharmacy care
- E- act with honesty, without deception

Answer:

C

Question Type: MultipleChoice

Rx Timolol 0.25% dropsMitte: 15 mLSig: gtt. i o.d. BIDOn the prescription label the instructions to the patient should read:

Options:

- A- apply one drop into both eyes twice daily.
- B- instil one drop into the left eye twice daily.
- **C-** instil one drop into the right eye twice daily.
- D- shake well and instil one drop into the left eye twice daily.
- E- shake well and instil one drop into the right ear twice daily.

Answer:

С

Question 8

Question Type: MultipleChoice

Factors that determine bioalence of two brands of a drug include:
I the taste of the preparations
Il the physical appearance of the preparations
III pharmacokinetic parameters of the preparations
Options:
A- I only
B- III only
C- I and II only
D- II and III only
E- I, II and III
Answer:
В
Question 9

Question Type:	MultipleChoice
-----------------------	----------------

A patient purchasing sublingual nitroglycerin tablets should be told to store the medication:

Options:

- A- in the refrigerator
- B- in a plastic vial with a childproof cap
- **C-** with the original cotton
- D- in an amber glass bottle with a metal cap
- E- in a warm, dry place

Answer:

D

Question 10

Question Type: MultipleChoice

Auxiliary labels which should be used on cotrimoxazole suspension include which of the following?

I Shake well before using
Il Take with plenty of fluids
III Exposure to sunlight may cause adverse reactions
Options:
A- I only
B- III only
C- I and II only
D- II and III only
E- I, II and III
Answer:
E
Question 11
Question Type: MultipleChoice

The pharmacist fills a prescription for sumatriptan 100 mg tablets for a migraine patient. Appropriate information to provide to the patient includes which of the following?

Options:

- A- If the sumatriptan does not relieve the headache within four hours, ergotamine may be used.
- B- If no relief is achieved in two hours, sumatriptan may be repeated.
- C- If the headache is relieved but another headache occurs eight hours later, sumatriptan may be used for the second headache.
- **D-** The maximum dosage in any 24 hour period is six tablets.
- E- If relief is not achieved, no other medication can be used for 24 hours.

Answer:

C

Question 12

Question Type: MultipleChoice

In an adequately powered, randomized controlled trial conducted over 3 years, a specific serious side effect (i.e. reduction in leukocytes) with conventional therapy is seen in 0.5% of the study sample. In patients who receive a newly discovered drug, only 0.45% experience the same side effect. Based on these results, the minimum number of patients that would have to receive the new drug for 3 years to statistically demonstrate the prevention of one episode of this side effect in at least one patient is:

0	pti	0	ns	
	Ρ	•		-

A- 15.

B- 20.

C- 150.

D- 200.

E- 2000.

Answer:

Е

To Get Premium Files for FPGEE Visit

https://www.p2pexams.com/products/fpgee

For More Free Questions Visit

https://www.p2pexams.com/nabp/pdf/fpgee

